

How To Treat An N -Level System : A Proposal *

Kazuyuki FUJII [†]

Department of Mathematical Sciences

Yokohama City University

Yokohama, 236-0027

Japan

Abstract

In this paper we propose a Hamiltonian of the n -level system by making use of generalized Pauli matrices.

An atom has usually many (finite or infinite) energy levels. However, to treat infinitely many ones at the same time is not realistic, so we treat an atom with finite energy levels. In the following we would like to consider an atom with n energy levels interacting with external (periodic) fields like $\cos(\omega t + \phi)$ (an n -level system in our terminology).

In Quantum Optics it is standard to consider a 2-level system, see for example [1] and [2]. The n -level system might be considered as a composite of 2-level systems. As for recent developments of the 2-level system including constructions of solutions in the strong coupling regime, see [3], [4], [5], [6], [13] and their references. See also [8] as a general introduction to two-level system, and [7], [10] as some applications to Quantum Optics. However we would like to treat an n -level system in a direct manner. Then we meet some troubles. First what is a Hamiltonian describing such a system ? As far as we

*This is prepared for the coming 11-th Numazu Meeting (6-8/March/2003)

[†]E-mail address : fujii@yokohama-cu.ac.jp

know such a Hamiltonian has not been known. As for some examples of n -level system see [12], [17], [18] and their references.

In this paper we propose such a Hamiltonian. In the 2-level system the Hamiltonian is written by making use of Pauli matrices $\{\sigma_1, \sigma_3\}$ (see the references above), so in the n -level system the Hamiltonian should be given by making use of generalized Pauli matrices $\{\Sigma_1, \Sigma_3\}$. We will write down it. We believe that it is a natural generalization

Our real aim of this study is to apply some developments in this paper to the theory of qudits in Quantum Computation, see for example [11] and [15]. In it we must first of all construct all unitary elements in $U(n)$ (a universality of 1-qudit).

We expect that this work will become a starting point toward this direction ¹.

Let $\{\sigma_1, \sigma_2, \sigma_3\}$ be Pauli matrices and $\mathbf{1}_2$ a unit matrix :

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{1}_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (1)$$

and $\sigma_+ = (1/2)(\sigma_1 + i\sigma_2)$, $\sigma_- = (1/2)(\sigma_1 - i\sigma_2)$.

First list the well-known properties of σ_1 and σ_3 :

$$\sigma_1^2 = \sigma_3^2 = \mathbf{1}_2, \quad \sigma_1^\dagger = \sigma_1, \quad \sigma_3^\dagger = \sigma_3, \quad \sigma_3\sigma_1 = -\sigma_1\sigma_3 = e^{\pi i}\sigma_1\sigma_3. \quad (2)$$

Let W be the Walsh-Hadamard matrix

$$W = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = W^{-1}, \quad (3)$$

then we can diagonalize σ_1 as $\sigma_1 = W\sigma_3W^{-1}$ by making use of this W .

Let us consider an atom with 2 energy levels E_0 and E_1 . Its Hamiltonian is in the diagonal form given as

$$H_0 = \begin{pmatrix} E_0 & 0 \\ 0 & E_1 \end{pmatrix}. \quad (4)$$

¹This paper may be deeply connected to [17] in Quantum Information Theory.

This is rewritten as

$$H_0 = \frac{E_0 + E_1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{E_0 - E_1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = \Delta_0 \mathbf{1}_2 + \Delta_1 \sigma_3, \quad (5)$$

where $\Delta_0 = (1/2)(E_0 + E_1)$ and $\Delta_1 = (1/2)(E_0 - E_1)$. Since we usually take no interest in constant terms, we set

$$H_0 = \Delta_1 \sigma_3, \quad \text{where} \quad \Delta_1 = (1/2)(E_0 - E_1). \quad (6)$$

Next we would like to extend 2-level system to general n -level system. For that we consider the 3-level one in detail as a test case. The Hamiltonian with three energy levels E_0, E_1, E_2 is

$$H_0 = \begin{pmatrix} E_0 & & \\ & E_1 & \\ & & E_2 \end{pmatrix}. \quad (7)$$

We want to rewrite this as (5). Let σ be $\exp(\frac{2\pi i}{3})$, then it is easy to see

$$\sigma^3 = 1, \quad \bar{\sigma} = \sigma^2, \quad 1 + \sigma + \sigma^2 = 0. \quad (8)$$

Since the matrix corresponding to σ_3 is considered as

$$\Sigma_3 = \begin{pmatrix} 1 & & \\ & \sigma & \\ & & \sigma^2 \end{pmatrix}, \quad (9)$$

we solve the equation

$$H_0 = \Delta_0 \mathbf{1}_3 + \Delta_1 \Sigma_3 + \Delta_2 \Sigma_3^2 \quad (10)$$

with indeterminants $\Delta_0, \Delta_1, \Delta_2$. Then we have

$$\begin{cases} \Delta_0 + \Delta_1 + \Delta_2 = E_0 \\ \Delta_0 + \sigma \Delta_1 + \sigma^2 \Delta_2 = E_1 \\ \Delta_0 + \sigma^2 \Delta_1 + \sigma \Delta_2 = E_2 \end{cases} \iff \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma & \sigma^2 \\ 1 & \sigma^2 & \sigma \end{pmatrix} \begin{pmatrix} \Delta_0 \\ \Delta_1 \\ \Delta_2 \end{pmatrix} = \begin{pmatrix} E_0 \\ E_1 \\ E_2 \end{pmatrix}, \quad (11)$$

so we obtain

$$\begin{pmatrix} \Delta_0 \\ \Delta_1 \\ \Delta_2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma & \sigma^2 \\ 1 & \sigma^2 & \sigma \end{pmatrix}^{-1} \begin{pmatrix} E_0 \\ E_1 \\ E_2 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma^2 & \sigma \\ 1 & \sigma & \sigma^2 \end{pmatrix} \begin{pmatrix} E_0 \\ E_1 \\ E_2 \end{pmatrix}, \quad (12)$$

or

$$\begin{cases} \Delta_0 = \frac{1}{3}(E_0 + E_1 + E_2), \\ \Delta_1 = \frac{1}{3}(E_0 + \sigma^2 E_1 + \sigma E_2), \\ \Delta_2 = \frac{1}{3}(E_0 + \sigma E_1 + \sigma^2 E_2). \end{cases} \quad (13)$$

Therefore the Hamiltonian which neglects the constant term is written as

$$H_0 = \Delta_1 \Sigma_3 + \Delta_2 \Sigma_3^2 = \Delta_1 \Sigma_3 + \bar{\Delta}_1 \Sigma_3^\dagger. \quad (14)$$

These Δ_1 and Δ_2 may be considered as a complex "energy difference" among E_0, E_1, E_2 .

By the way, the matrix in the right hand side of (12) (replaced $1/3$ with $1/\sqrt{3}$)

$$W = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma^2 & \sigma \\ 1 & \sigma & \sigma^2 \end{pmatrix} \in U(3) \quad (15)$$

is very familiar to us. Let Σ_1 and Σ_3 be generators of generalized Pauli matrices in the case of $n = 3$, namely

$$\Sigma_1 = \begin{pmatrix} 0 & & 1 \\ 1 & 0 & \\ & 1 & 0 \end{pmatrix}, \quad \Sigma_3 = \begin{pmatrix} 1 & & \\ & \sigma & \\ & & \sigma^2 \end{pmatrix}. \quad (16)$$

Then it is easy to see

$$\Sigma_1^3 = \Sigma_3^3 = \mathbf{1}_3, \quad \Sigma_1^\dagger = \Sigma_1^2, \quad \Sigma_3^\dagger = \Sigma_3^2, \quad \Sigma_3 \Sigma_1 = \sigma \Sigma_1 \Sigma_3. \quad (17)$$

Now we can show that Σ_1 can be diagonalized by making use of W above

$$\Sigma_1 = W \Sigma_3 W^\dagger = W \Sigma_3 W^{-1}. \quad (18)$$

In fact

$$\begin{aligned}
W\Sigma_3W^\dagger &= \frac{1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma^2 & \sigma \\ 1 & \sigma & \sigma^2 \end{pmatrix} \begin{pmatrix} 1 & & \\ & \sigma & \\ & & \sigma^2 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma & \sigma^2 \\ 1 & \sigma^2 & \sigma \end{pmatrix} \\
&= \frac{1}{3} \begin{pmatrix} 1 & \sigma & \sigma^2 \\ 1 & 1 & 1 \\ 1 & \sigma^2 & \sigma \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \sigma & \sigma^2 \\ 1 & \sigma^2 & \sigma \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 0 & 0 & 3 \\ 3 & 0 & 0 \\ 0 & 3 & 0 \end{pmatrix} = \Sigma_1,
\end{aligned}$$

where we have used the relations of σ in (8).

A comment is in order. Since W corresponds to the Walsh–Hadamard matrix (3), so it may be possible to call W the generalized Walsh–Hadamard matrix.

Let us proceed with general case. The Hamiltonian with n energy levels $E_0, E_1, E_2, \dots, E_{n-1}$ is

$$H_0 = \begin{pmatrix} E_0 & & & & \\ & E_1 & & & \\ & & E_2 & & \\ & & & \ddots & \\ & & & & \ddots \\ & & & & & E_{n-1} \end{pmatrix}. \quad (19)$$

From the discussion in the case of $n = 3$ it is easy to rewrite the Hamiltonian. Let $\{\Sigma_1, \Sigma_3\}$ be generalized Pauli matrices (see [9], Appendix B)

$$\Sigma_1 = \begin{pmatrix} 0 & & & & 1 \\ 1 & 0 & & & \\ & 1 & 0 & & \\ & & \ddots & \ddots & \\ & & & \ddots & \ddots \\ & & & & 1 & 0 \end{pmatrix}, \quad \Sigma_3 = \begin{pmatrix} 1 & & & & \\ & \sigma & & & \\ & & \sigma^2 & & \\ & & & \ddots & \\ & & & & \ddots \\ & & & & & \sigma^{n-1} \end{pmatrix} \quad (20)$$

where σ is a primitive element $\sigma = \exp(\frac{2\pi i}{n})$ which satisfies

$$\sigma^n = 1, \quad \bar{\sigma} = \sigma^{n-1}, \quad 1 + \sigma + \dots + \sigma^{n-1} = 0. \quad (21)$$

Then it is easy to see

$$\Sigma_1^n = \Sigma_3^n = \mathbf{1}_n, \quad \Sigma_1^\dagger = \Sigma_1^{n-1}, \quad \Sigma_3^\dagger = \Sigma_3^{n-1}, \quad \Sigma_3 \Sigma_1 = \sigma \Sigma_1 \Sigma_3. \quad (22)$$

If we define a Vandermonde matrix W based on σ as

$$W = \frac{1}{\sqrt{n}} \begin{pmatrix} 1 & 1 & 1 & \cdot & \cdot & \cdot & 1 \\ 1 & \sigma^{n-1} & \sigma^{2(n-1)} & \cdot & \cdot & \cdot & \sigma^{(n-1)^2} \\ 1 & \sigma^{n-2} & \sigma^{2(n-2)} & \cdot & \cdot & \cdot & \sigma^{(n-1)(n-2)} \\ \cdot & \cdot & \cdot & & & & \cdot \\ \cdot & \cdot & \cdot & & & & \cdot \\ 1 & \sigma^2 & \sigma^4 & \cdot & \cdot & \cdot & \sigma^{2(n-1)} \\ 1 & \sigma & \sigma^2 & \cdot & \cdot & \cdot & \sigma^{n-1} \end{pmatrix}, \quad (23)$$

then it is not difficult to see

$$\Sigma_1 = W \Sigma_3 W^\dagger = W \Sigma_3 W^{-1}. \quad (24)$$

That is, Σ_1 can be diagonalized by making use of W .

We set for simplicity

$$\mathbf{\Delta} = (\Delta_0, \Delta_1, \dots, \Delta_{n-2}, \Delta_{n-1})^t, \quad \mathbf{E} = (E_0, E_1, \dots, E_{n-2}, E_{n-1})^t, \quad (25)$$

and the relation like (12) between them is given by

$$\mathbf{\Delta} = \frac{1}{\sqrt{n}} W \mathbf{E}, \quad (26)$$

or explicitly

$$\Delta_j = (1/n) \left(E_0 + \sigma^{n-j} E_1 + \sigma^{2(n-j)} E_2 + \dots + \sigma^{(n-1)(n-j)} E_{n-1} \right)$$

for $0 \leq j \leq n-1$. Then the Hamiltonian which neglects the constant term is written as

$$H_0 = \sum_{j=1}^{n-1} \Delta_j \Sigma_3^j. \quad (27)$$

Next let us incorporate an interaction into the model. We consider an atom with two energy levels which interacts with external (periodic) field with $g\cos(\omega t)$. The Hamiltonian in the dipole approximation is given by

$$H = H_0 + g \cos(\omega t)\sigma_1 = \Delta_1\sigma_3 + g \cos(\omega t)\sigma_1, \quad \cos(\omega t) = (1/2)(e^{i\omega t} + e^{-i\omega t}), \quad (28)$$

where Δ_1 is the constant in (6), ω the frequency of the external field, g the coupling constant between the external field and the atom. In the following we cannot assume the rotating wave approximation (which neglects the fast oscillating terms), namely the Hamiltonian given by

$$H = \Delta_1\sigma_3 + \frac{g}{2} (e^{i\omega t}\sigma_+ + e^{-i\omega t}\sigma_-). \quad (29)$$

We would like to extend this to the general case. We again note that Σ_1 is not hermitian, so we must take this fact into consideration. The interaction term corresponding to $\cos(\omega t)\sigma_1$ is considered to be

$$\frac{1}{2} (e^{i\omega t}\Sigma_1 + e^{-i\omega t}\Sigma_1^\dagger) \quad (30)$$

near to the above (29). Let us explain the reason. The diagonalized form of $\cos(\omega t)\sigma_1$ is given by

$$\cos(\omega t)\sigma_1 = W \begin{pmatrix} \cos(\omega t) & \\ & -\cos(\omega t) \end{pmatrix} W^{-1} = W \begin{pmatrix} \cos(\omega t) & \\ & \cos(\omega t + \pi) \end{pmatrix} W^{-1}$$

by making use of W in (3). On the other hand it is easy to see

$$\frac{1}{2} (e^{i\omega t}\Sigma_1 + e^{-i\omega t}\Sigma_1^\dagger) = W D W^{-1} \quad (31)$$

by making use of W in (23), where D is a diagonal matrix

$$D = \begin{pmatrix} \cos(\omega t) & & & & \\ & \cos(\omega t + 2\pi/n) & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \cos(\omega t + 2\pi(n-1)/n) \end{pmatrix}. \quad (32)$$

Therefore this is a natural generalization of $\cos(\omega t)\sigma_1$.

Now we are in a position to state our proposal :

Proposal The Hamiltonian in the general case is

$$H = H_0 + \frac{g}{2} \left(e^{i\omega t} \Sigma_1 + e^{-i\omega t} \Sigma_1^\dagger \right) = \sum_{j=1}^{n-1} \Delta_j \Sigma_3^j + \frac{g}{2} \left(e^{i\omega t} \Sigma_1 + e^{-i\omega t} \Sigma_1^\dagger \right) \quad (33)$$

where g is a coupling constant of the system. In particular in the case of $n = 3$

$$H = \Delta_1 \Sigma_3 + \bar{\Delta}_1 \Sigma_3^\dagger + \frac{g}{2} \left(e^{i\omega t} \Sigma_1 + e^{-i\omega t} \Sigma_1^\dagger \right) \quad (34)$$

by (14).

We have written down the Hamiltonian in the general case in terms of generalized Pauli matrices, however don't know whether or not it is actually reasonable in Quantum Optics or Condensed Matter Physics (refer to [16]). Further considerations will be needed.

We conclude this paper by stating that constructions of solutions under some ansatz (in the weak or strong coupling regime) are not difficult and will be published in another paper, [14].

References

- [1] L. Allen and J. H. Eberly : Optical Resonance and Two-Level Atoms, Wiley, New York, 1975.
- [2] A. Corney : Atomic and Laser Spectroscopy, Oxford University Press, 1977.
- [3] C. E. Creffield : Location of crossing in the spectrum of a driven two-level system, cond-mat/0301168.
- [4] J. C. A. Barata and W. F. Wreszinski : Strong Coupling Theory of Two Level Atoms in Periodic Fields, Phys. Rev. Lett. 84(2000), 2112, physics/9906029.
- [5] A. Santana, J. M. Gomez Llorente and V. Delgado : Semiclassical dressed states of two-level quantum systems driven by nonresonant and/or strong laser fields, J. Phys. B 34(2001), 2371, quant-ph/0011015.

- [6] M. Frasca : Theory of dressed states in quantum optics, Phys. Rev. A 60(1999), 573, quant-ph/9811037.
- [7] M. Frasca : Rabi oscillations and macroscopic quantum superposition states, Phys. Rev. A 66(2002), 023810, quant-ph/0111134.
- [8] M. Frasca : A modern review of the two-level approximation, quant-ph/0209056.
- [9] K. Fujii : Introduction to Grassmann Manifolds and Quantum Computation, J. Applied Math, 2(2002), 371, quant-ph/0103011.
- [10] K. Fujii : Mathematical Structure of Rabi Oscillations in the Strong Coupling Regime, to appear in J. Phys. A, quant-ph/0203135.
- [11] K. Fujii : Exchange Gate on the Qudit Space and Fock Space, submitted, quant-ph/0207002.
- [12] K. Fujii : N-Level System Interacting with Single Radiation Mode and Multi Cat States of Schrödinger in the Strong Coupling Regime, quant-ph/0210166.
- [13] K. Fujii : Two-Level System and Some Approximate Solutions in the Strong Coupling Regime, quant-ph/0301145.
- [14] K. Fujii et al : N-Level System and Some Approximate Solutions in the Strong Coupling Regime (tentative), in preparation.
- [15] K. Funahashi : The Controlled-U and unitary transformations on two qudits, in preparation.
- [16] P. Meystre and M. Sargent III : Elements of Quantum Optics, Springer-Verlag, 1990.
- [17] M. Zukowski, A. Zeilinger and M. A. Horne : Realizable higher-dimensional two-particle entanglements via multiport beam splitters, Phys. Rev. A 55(1997), 2564.

- [18] D. Ellinas and E. G. Floratos : Prime decomposition and correlation measure of finite quantum systems, J. Phys. A 32(1999), L63, quant-ph/9806007.